

Chalek Juice: An alternative diet therapy of diabetes mellitus

by Erma Nurjanah Widiastuti


Submission date: 30-Jan-2023 03:47PM (UTC+0700)

Submission ID: 2002381564

File name: PUBLIKASI_JUS_CHALEK_939-Article_Text-3102-1-10-20221127.pdf (895.84K)

Word count: 1208

Character count: 6826



Open Access

Photo Essay

POLITEKNIK KESEHATAN KEMENKES PALANGKA RAYA: HEALTH FORUM AND INTERNATIONAL SEMINAR
THE NEW NORMAL : Creating a Pleasant Virtual Communication

Chalek Juice: An alternative diet therapy of diabetes mellitus

Vinna Maulidha Yunianitamaras*, Erma Nurjanah Widiastuti, Cucu Rahayu, Agnescia Clarissa Sera

Department of Nutrition, Poltekkes Kemenkes Palangka Raya, Indonesia

*Corresponding author's email: vinnamaulidhay@gmail.com

DOI: [10.35898/ghmj-52939](https://doi.org/10.35898/ghmj-52939)

⁵ Selection and peer-review under responsibility of the scientific committee and the editorial board of the Annual Health Forum and International Seminar of the Politeknik Kesehatan Kemenkes Palangka Raya

¹ © Yayasan Aliansi Cendekiawan Indonesia Thailand (Indonesian Scholars' Alliance). This is an open-access following Creative Commons License Deed - Attribution-NonCommercial-ShareAlike 4.0 International (CC BY-NC-SA 4.0)

Diabetes mellitus (DM) is ⁷ chronic disease caused by insufficient amounts of insulin due to the malfunction of pancreatic organ (Ministry of Health of Republic of Indonesia, 2019; Arsono, 2005). It will impact productivity, thereby reducing the quality of human resources. Indonesia is among the 10 largest countries in the world with DM cases and is ranked 4th (Damayanti, 2015; Ministry of Health of Republic of Indonesia, 2018; Irawan, 2010). Health treatment for DM requires expensive costs, especially for patients with complications. This ² encourages some people to look for herbal therapy, such as bitter melon (*Momordica charantia L*) (Ministry of Health of Republic of Indonesia, 2014; Katmo and Pramono, 2006).



Figure 1 Bitter Melon (*Momordica charantia L*)

Shown in Figure 1, Bitter Melon or known in Indonesia as Pare, contains saponins, flavonoids, polyphenols, and alkaloids, which may improve pancreatic beta cells, thereby increasing insulin production (Suiraoaka, 2012). Besides, bitter melon also contains charantin, an active antidiabetic substance that is believed to reduce blood ⁶ sugar levels in the body. The decrease in blood sugar level is marked by lectin, carbohydrate-binding proteins expressed on the surface of the body's cells (Martínez-Alarcón et al., 2018; Tandra, 2013). This prompted the authors to create an alternative diet therapy formula named Chalek Juice, which stands for Charantin and Lectin.



Figure 2. Chalek Juice made from a mixture of bitter melon, cucumber, green apple and lemon)

As the name implies, bitter melon is dominated by bitter taste. It is mixed with cucumber, green apple and lemon to get a fresh and less bitter taste. To make Chalek Juice, there are six steps. First, wash the fruits and vegetables followed by cutting them into small pieces. To reduce the bitterness, soak the bitter melon into salt water for about 10 minutes (Setiawan, 1993; Rukmana, 2006), as shown in Figure 3.



Figure 3 Soaking process of bitter melon into salt water to reduce the bitterness

The bitter melon, cucumber and green apple are crushed separately using a food processor. Before mixing all the ingredients, bitter melon, cucumber and green apple are strained from their pulps. To get a glass of Chalek Juice (250 mL), mix 120 mL of bitter melon, 60 mL of green apple, 60 mL of cucumber and 20 mL of lemon juice, and mix it (Figure 4). Finally, the Chalek Juice is ready to drink. It is highly recommended to drink it at night before going to bed.



Figure 4. Mixing all ingredients

A glass of Chalek juice contains 106 calories, 19.6 grams of carbohydrates, 4.1 grams of protein, 2.9 grams of fat, 2.4 grams of fiber, 60 mg of calcium and 42 mg of sodium. All ingredients used in the juice making are affordable and very easy to find at markets or convenient stores.



Figure 5. Blood sugar level testing after two hours of consuming chalek juice

In addition, bitter melon likely increases beta-cells or cells in the pancreas that function to produce insulin in the pancreas, and it is believed to help insulin secretion. This makes bitter melon often associated with diabetes (Fuangchan *et al.*, 2011). As shown in Figure 5, we measured the blood sugar level in the participants after 2 hours consuming Chalek Juice. Hasibuan and Manurung (2020) state that bitter melon juice was significantly effective in reducing blood sugar levels in patients with Type II diabetes mellitus ($p = 0.000$). However, a research by Gardiarini *et al.* (2018) using the hedonic test of the taste panel found that most of the panelists favored and really favored the bitter melon food (sautéed bitter melon) provided by the hospital. It is suggested, this innovation still needs to be improved in the future, especially its durability. This study found Chalek Juice only lasts a maximum of one week at cold temperatures and 30 minutes at a room temperature.

Consent

The informants (identifiable) have given their consent for their pictures to be used in the publication of this research.

Conflict of Interest

None.

Acknowledgments

The authors would thank to the whole academic community of Poltekkes Kemenkes Palangka Raya who support the development of Chalek Juice.

References

- Arsono, S. (2005). *Diabetes Melitus Sebagai Faktor Risiko Kejadian Gagal Ginjal Terminal Tesis*. Semarang: Program Pasca Sarjana Universitas Diponegoro.
- Damayanti, S. (2015). *Diabetes Mellitus & Penatalaksanaan Keperawatan*. Yogyakarta: Nuha Medika
- Fuangchan, A., Sonthisombat, P., Seubnukarn, T., Chanouan, R., Chotchaisuwat, P., Sirigulsatien, V., Ingkaninan, K., Plianbangchang, P., & Haines, S. T. (2011). Hypoglycemic effect of bitter melon compared with metformin in newly diagnosed type 2 diabetes patients. *Journal of Ethnopharmacology*, 134(2), 422-428. <https://doi.org/10.1016/j.jep.2010.12.045>
- Gardiariini, P., Setyawati, R., Setyaningsih, A. (2018). Pengembangan Menu Berbahan Dasar Pare sebagai Menu Diet Khusus Penderita Diabetes Melitus di Rumah Sakit. *SNITT-Politeknik Negeri Balikpapan*. 263-268.
- Hasibuan, D.C., & Manurung, D.M. (2020). Efektifitas Pare Terhadap Penurunan Kadar Gula Darah Pada Penderita Diabetes Mellitus Tipe II. *Jurnal Kesehatan Ilmiah Indonesia*, 5 (2), 68-73. <http://dx.doi.org/10.51933/health.v5i2.318>
- Irawan, Dedi. 2010. *Prevalensi dan Faktor Resiko Kejadian Diabetes Mellitus Tipe 2 di daerah Urban Indonesia (Analisa Data Sekunder Riskesdas 2007)*. Thesis: Universitas Indonesia.
- Katmo dan Pramono, S. (2006). *Tingkat Manfaat Dan Keamanan Tumbuhan Obat dan Obat Tradisional*. Yogyakarta: Fakultas Farmasi UGM.
- Martínez-Alarcón, D., Blanco-Labra, A., & García-Gasca, T. (2018). Expression of Lectins in Heterologous Systems. *International journal of molecular sciences*, 19(2), 616. <https://doi.org/10.3390/ijms19020616>
- Ministry of Health of Republic of Indonesia. (2014). *Situasi Dan Analisis Diabetes*. Jakarta Selatan : Pusat Data dan Informasi.
- Ministry of Health of Republic of Indonesia. (2018). *Hasil Utama RISKESDAS Indonesia Tahun 2018*. Jakarta: Kemenkes RI.
- Ministry of Health of Republic of Indonesia. 2019. *Pusat Data dan Informasi Kementerian Kesehatan RI Hari Diabetes Sedunia Tahun 2018*. Jakarta: Kemenkes RI.
- Rukmana, R. (2006). *Budi Daya Pare*. Yogyakarta: Kanisius.
- Setiawan, Iwan Ade. (1993). *Pare dan Labu*. Jakarta: PT Penebar Swadaya.
- Suiraoaka, IP. (2012). *Penyakit Degeneratif. Mengenal, Mencegah dan Mengurangi Faktor Resiko 9 Penyakit Degeneratif*. Yogyakarta : Nuha Medika.
- Tandra, H. (2013). *Life Healthy With Diabetes (Diabetes Mengapa & Bagaimana?)*. Yogyakarta: Rapha PUBLISHING.

Cite this article as: Yunianitamara VM, Widiastuti EN, Rahayu C., Sera AC. Chalek Juice: An alternative diet therapy of diabetes mellitus. *GHMJ (Global Health Management Journal)*. 2022; 5(2):128-131. [doi:10.35898/GHMJ-52939](https://doi.org/10.35898/GHMJ-52939)

Chalek Juice: An alternative diet therapy of diabetes mellitus

ORIGINALITY REPORT

12%

SIMILARITY INDEX

7%

INTERNET SOURCES

6%

PUBLICATIONS

6%

STUDENT PAPERS

PRIMARY SOURCES

1	Submitted to Politeknik Kesehatan Kemenkes Semarang Student Paper	4%
2	www.scitepress.org Internet Source	2%
3	Submitted to Eastern Mennonite University Student Paper	2%
4	nusantarahasanajournal.com Internet Source	1%
5	www.elsevier.es Internet Source	1%
6	"Aquatic Lectins", Springer Science and Business Media LLC, 2022 Publication	1%
7	repo.stikesicme-jbg.ac.id Internet Source	1%

Exclude quotes

Off

Exclude matches

Off

Exclude bibliography On