



POLITEKNIK KESEHATAN KEMENKES PALANGKA RAYA: HEALTH FORUM AND INTERNATIONAL SEMINAR  
THE NEW NORMAL : Creating a Pleasant Virtual Communication

# Risk factors of stunting in toddlers aged 24-59 months living in floating houses on the Kahayan River of Palangka Raya City: A case-control study

Mars Khendra Kusfriyadi\*, Sugiyanto Sugiyanto

Department of Nutrition, Poltekkes Kemenkes Palangka Raya, Indonesia

\*Corresponding author's email: [marikhend@gmail.com](mailto:marikhend@gmail.com)

DOI: [10.35898/ghmj-51605](https://doi.org/10.35898/ghmj-51605)

---

*Selection and peer-review under responsibility of the scientific committee and the editorial board of the Annual Health Forum and International Seminar of the Politeknik Kesehatan Kemenkes Palangka Raya*

---

© Yayasan Aliansi Cendekiawan Indonesia Thailand (Indonesian Scholars' Alliance). This is an open-access following Creative Commons License Deed - Attribution-NonCommercial-ShareAlike 4.0 International (CC BY-NC-SA 4.0)

Central Kalimantan, one of the provinces in Indonesia, has many tributaries that flow into Kahayan River. The river has a length of 450 m<sup>2</sup> and a width of 100 m<sup>2</sup>. Around the river, we can see portraits of the people and their daily activities. Residential housing on Kahayan River is divided into two models, namely *rumah lanting* or floating house (raft house), and *rumah tiang* or pillar houses (Hamidah *at al.*, 2014).



*Figure 1 Kahayan Bridge Passing Over Kahayan River and A Row of Floating Houses*

There were 342 floating houses on the river. The people use the river's water for cooking, washing, drinking, bathing, and defecation. These activities may devastate the environmental quality around the river. Poor quality environmental quality will also impact the public health, especially children.

The prevalence of stunting in toddlers in Central Kalimantan exceeded the national average of 39.0% and in Palangka Raya City, it reached 29.7% (Dinas Kesehatan, 2018). This background suggest it already exceeded the standards of WHO as well (20%). Thus, it must be addressed promptly with proper actions. To solve public health problems, especially stunting, cooperation across sectors and research for spatially analyzing stunting risk factors are required. Community residential condition also warrants attention from the local authorities. To look further into the stunting issue, we studied stunting risk factors that may harm the community on the river.



*Figure 2 Activities Around Kahayan River*

This observational study used a case control design to find out risk factors of stunting in toddlers living in floating houses on Kahayan River in Palangka Raya. Eighty stunting toddlers aged between 24-59 months were studied. Forty toddlers who lived in floating houses were treated as a case group, and 40 toddlers who lived in Kahayan watersheds became a control group.



*Figure 3 Visiting Floating House Residence Approximately for 20-30 Minutes From The Port*



*Figure 4 Researcher and The Floating Houses*

Approximately 98% of the people lived permanently on the houses, and only 12 % lived less than a year.



*Figure 5 Floating Houses as Homes For Fish Farmers*



*Figure 6 Landscape of Floating Houses From The Boat*



*Figure 7 Floating Toilet on The River*

Stunting risk factors examined in this study include child, maternal, and environmental factors. The multivariate regression analysis showed the behavioral factor i.e., garbage disposal to the river was the most influential for causing stunting (OR = 0.210), followed by infectious disease factor such as acute respiratory tract infections which most frequently occurred (OR = 0.190), food intake (OR = 0.163), fruit consumption (OR = 0.029), mother's knowledge of nutrition (OR = 0.026) and education level of father (OR = 0.009).



*Figure 8 A Stunting Child with His Mother Who Had Lived For 20 Years in The Floating House*



*Figure 9 Measuring Toddler’s Height on Floating Houses*



*Figure 10 Researchers Also Measured The Height of Toddler Parents*

Toddlers living in floating houses were more likely to be at risk of stunting those living in the watersheds. We calculated the possibility of stunting occurrence (p) with the following formula:

$$p = \frac{\text{Exp} (5.496 + [(-1.562 \text{ GD}) + (-1.663 \text{ ID}) + (-1.817 \text{ FI}) + (-3.525 \text{ FC}) + (-3.638 \text{ MK}) + (-4.682 \text{ FE})])}{1 + \text{Exp} (5.496 + [(-1.562 \text{ GD}) + (-1.663 \text{ ID}) + (-1.817 \text{ FI}) + (-3.525 \text{ FC}) + (-3.638 \text{ MK}) + (-4.682 \text{ FE})])}$$

GD = garbage disposal into the river, ID = infectious disease factor, FI = Food intake, FC = fruit consumption, MK = Mother’s knowledge about nutrition, FE = level education of father

This model that shows stunting risk factors aligns with the models created by UNICEF and UN and specifically portrays stunting prevalence on Kahayan River.



*Figure 11 Research Data Collection by Enumerators*



*Figure 12 Habit of Dumping Garbage Into Rivers*

Local government, especially the Palangkaraya District Health Office is able to use results of this study for better policy and intervention planning to improve the nutritional status of the community around Kahayan River. Besides, our research gives implications for tourist area development around the river.

### **Consent**

Participants of the research have given their consent for use of their photographs as needed in this study. All of the photographs belong to the researchers under this study.

### **Conflict of Interest**

None.

### **Acknowledgments**

This study was funded by the Health Polytechnic of Palangka Raya in 2018.

### **References**

Bunga Ch Rosha, Hardinsyah Dan Yayuk Farida Baliwati. 2012. Analisis Determinan Stunting Anak 0-23 Bulan Pada Daerah Miskin Di Jawa Tengah Dan Jawa Timur Penel Gizi Makan. *The Journal Nutrition And Food Research*, 35(1):34-41. Retrieved from: <https://ejournal.litbang.kemkes.go.id/index.php/pgm/article/view/3081>

- Departemen Kesehatan. 2005. *Rencana Aksi Nasional Pencegahan dan Penanggulangan Gizi Buruk 2005-2009*. Jakarta : Depkes RI, Direktorat Jenderal Pelayanan Kesehatan.
- Dinas Kesehatan Provinsi Kalimantan Tengah. 2018. Laporan Pemantauan Status Gizi Provinsi Tahun 2017. Dian Hani Ulfani, Drajat Martianto, Dan Yayuk Farida Baliwati. 2011. Faktor-Faktor Sosial Ekonomi Dan Kesehatan Masyarakat Kaitannya Dengan Masalah Gizi Underweight, Stunted, Dan Wasted Di Indonesia: Pendekatan Ekologi Gizi. *Journal of Nutrition and Food*, 6(1): 59-65. DOI: <https://doi.org/10.25182/jgp.2011.6.1.59-65>
- Erna Kusumawati, Setiyowati Rahardjo, Hesti Permata Sari. 2015. Model Pengendalian Faktor Risiko Stunting Pada Anak Usia Di Bawah Tiga Tahun. *Jurnal Kesehatan Masyarakat Nasional* , 9(3). Retrieved from. <http://dx.doi.org/10.21109/kesmas.v9i3.572>
- Firmanu Cahyono, Stefanus Pieter Manongga, Intje Picauly. 2016. Faktor Penentu Stunting Anak Balita pada Berbagai Zona Ekosistem di Kabupaten Kupang *Jurnal Gizi Pangan*, 11(1): 9-18. DOI: <https://doi.org/10.25182/jgp.2016.11.1.%25p>
- Noor Hamidah, R. Rijanta Bakti Setiawan Muh. Aris Marfai. 2014. Kajian Transportasi Sungai Untuk Menghidupkan Kawasan Tepian Sungai Kahayan Kota Palangkaraya. *Jurnal Tata Loka*, 16(1),1-17. Retrieved from: <https://www.academia.edu/48513047/>
- Novrianti. 2016. Pengaruh Aktivitas Masyarakat di pinggir Sungai (Rumah Terapung) terhadap Pencemaran Lingkungan Sungai Kahayan Kota Palangka Raya Kalimantan Tengah. *Media Ilmiah Teknik Lingkungan*, 1(2), 35-39. Retrieved from: <https://www.neliti.com/publications/258536/>
- Suprptini Dan Dwi Hapsari. 2011. Status Gizi Balita Berdasarkan Kondisi Lingkungan Dan Status Ekonomi (Data Riskesdas 2007). *Jurnal Ekologi Kesehatan*, 10 (2), 103-113. Retrieved from: <https://docobook.com/>

**Cite this article as:**

Kusfriyafi MK, Sugiyanto S. Risk factors of stunting in toddlers aged 24-59 months living in floating houses on the Kahayan River of Palangka Raya City : A case-control study. *GHMJ (Global Health Management Journal)*. 2022; 5(1):81-87. [doi:10.35898/ghmj-51605](https://doi.org/10.35898/ghmj-51605)